

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 633 022 A2

(12)

EUROPEAN PATENT APPLICATION(21) Application number: **94109872.5**(51) Int. Cl.⁸: **A61K 31/365, A61K 31/70**(22) Date of filing: **27.06.94**(30) Priority: **09.07.93 JP 194182/93**(43) Date of publication of application:
11.01.95 Bulletin 95/02(84) Designated Contracting States:
CH DE FR GB IT LI SE

(71) Applicant: **KUREHA CHEMICAL INDUSTRY CO., LTD.**
9-11, Horidome-cho, 1-chome
Nihonbashi
Chuo-ku
Tokyo 103 (JP)


(72) Inventor: **Watanabe, Koju**
2-5-7, Tsurumai
Sakado-shi,

Saitama 350-02 (JP)**Inventor: Nilmura, Koichi****Rune Warabi 1-718,****1-17-30, Chuo****Warabi-shi,****Saitama 335 (JP)****Inventor: Umekawa, Kiyonori****5-4-309, Mihama****Urayasu-shi,****Chiba 279 (JP)**

(74) Representative: **Minderop, Ralph H. Dr. rer.nat.**
et al

Cohausz & Florack**Patentanwälte****Bergiusstrasse 2 b****D-30655 Hannover (DE)**(54) **Chondroprotective agents.**

(57) A chondroprotective agent comprising a flavonoid compound of the general formula (I):



wherein R¹ to R⁹ are, independently, a hydrogen atom, hydroxyl group, or methoxyl group and X is a single bond or a double bond, or a stereoisomer thereof, or a naturally occurring glycoside thereof is disclosed. The above compound strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage, and thus, is extremely effective for the treatment of arthropathy.

BACKGROUND OF THE INVENTION

1. Field of the Invention

5 The present invention relates to an agent for protecting cartilage, i.e., a chondroprotective agent, more particularly, a chondroprotective agent containing a flavonoid compound or a stereoisomer thereof, or a naturally occurring glycoside thereof.

2. Description of the Related Art

10

There are various types of arthropathy, for example, rheumatoid arthritis, rheumatic fever, and osteoarthritis. Many people particularly suffer from rheumatoid arthritis and osteoarthritis. These diseases have been studied as the major types of arthropathy. There are congenital and secondary osteoarthritis, and further primary osteoarthritis caused by degeneration of the articular cartilage along with aging. Patients
15 suffering from primary osteoarthritis have recently been increasing along with the increase in the population of the aged.

Although there are considerable differences of the causes and conditions between rheumatoid arthritis and osteoarthritis, the articular function becomes eventually obstructed by the destruction of the cartilage in both of rheumatoid arthritis and osteoarthritis.

20 The first choice of medicines for the treatment of rheumatic diseases such as rheumatoid arthritis, rheumatic fever, systemic lupus erythematosus, and osteoarthritis are analgesic and anti-inflammatory agents, for example, aspirin or indometacin. Further, gold compounds such as Shiosol, immunomodulators, steroids, or D-penicillamine are used as medicines for the treatment of rheumatoid arthritis.

The above conventional analgesic and anti-inflammatory agents, however, were not effective against the
25 destruction of the articular cartilage, and in fact, sometimes exhibited adverse effect in experiments using chondrocytes. Further, no inhibitory effect on articular cartilage destruction was also observed in the above-mentioned medicines for the treatment of rheumatoid arthritis.

It is known that flavonoids may be used as an agent for protecting a blood vessel and further in the following pharmaceutical applications: a virus genome deactivating agent for apigenin, chrysin, morin, fisetin, and baicalein [Japanese Unexamined Patent Publication (Kokai) No. 2-101013], an agent for
30 determining the function of polymorphonuclear leukocyte for flavonoids [Japanese Unexamined Patent Publication (Kokai) No. 63-253254], an oral agent for suppressing smoking for flavonoids [Japanese Unexamined Patent Publication (Kokai) No. 4-46119], treatment of high protein edema for rutin, diosmin, and the like (U.S. Patent No. 5,096,887), an anti-tumor agent containing flavonoids [Japanese Unexamined
35 Patent Publication (Kokai) No. 3-275625], an anti-tumor agent containing apigenin [Japanese Examined Patent Publication (Kokoku) No. 3-61644], an agent for suppressing the formation of peroxy lipid for hesperetin, kaempferol, and the like [Japanese Unexamined Patent Publication (Kokai) No. 3-5423], an anti-tumor agent containing kaempferol [Japanese Unexamined Patent Publications (Kokai) No. 4-103529 and
40 No. 4-103532], a calcium antagonist for hesperidin and luteolin [Japanese Unexamined Patent Publication (Kokai) No. 4-243822], a sialidase inhibitor for luteolin [Japanese Unexamined Patent Publication (Kokai) No. 64-42427], an anti-retrovirus agent for luteolin [Japanese Unexamined Patent Publication (Kokai) No. 3-7224], an anti-HBV (hepatitis B virus) agent for quercetin [Japanese Unexamined Patent Publication (Kokai) No. 4-234320], and the like.

Flavonoids have not, however, been known to be useful as chondroprotective agents.

45


SUMMARY OF THE INVENTION

The present inventors engaged in intensive research to develop a chondroprotective agent for suppressing the destruction of the articular cartilage and as a result found that the particular flavonoid
50 compounds and stereoisomers thereof, and the naturally occurring known glycosides thereof showed significant inhibition of the depletion of proteoglycan which is a major component of the cartilage matrix, and therefore, are useful as a chondroprotective agent for prohibiting the destruction of the articular cartilage.

Accordingly, the object of the present invention is to provide a chondroprotective agent containing as an
55 active ingredient a particular flavonoid compound or a stereoisomer thereof, or a naturally occurring known glycoside thereof.

Other objects and effects of the present invention will be clear from the following description.

The present invention relates to a chondroprotective agent comprising a flavonoid compound of the general formula (I):



wherein R¹ to R⁹ are, independently, a hydrogen atom, hydroxyl group, or methoxyl group and X is a single bond or a double bond, or a stereoisomer thereof, or a naturally occurring glycoside thereof (hereinafter referred to as "the present substance").

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The active ingredient of the chondroprotective agent according to the present invention is a flavonoid, which is widely present in the vegetable kingdom. Typical flavonoid compounds include flavones, flavonols, flavanones, and flavanonols. Flavanones contain an asymmetric carbon atom at the 2-position, and flavanonols contain asymmetric carbon atoms at the 2- and 3-positions, and such compounds may be present as the stereoisomers. These stereoisomers can also be used in the present invention. Further, the saccharides present in the above naturally occurring glycosides are not particularly limited. As examples of the naturally occurring glycosides, there may be mentioned glucoside, galactoside, fructoside, rhamnoside, rutinoid (that is, rhamnoglucoside), arabinoside, xyloside, apioglucoside, and robinobioside.

In the present invention, any naturally occurring flavonoids may be used as the above present substance. The flavonoid compounds and naturally occurring glycosides thereof shown in the following Table 1 are preferable.

Table 1

	No. Name	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹	X
5	1 Flavone	H	H	H	H	H	H	H	H	H	Double
	2. Chrysin	H	OH	H	OH	H	H	H	H	H	ditto
10	3. Toringin	H	OH	H	OGlu	H	H	H	H	H	ditto
	4. Primetin	OH	H	H	OH	H	H	H	H	H	ditto
	5 Apigenin	H	OH	H	OH	H	H	H	OH	H	ditto
	6 Cosmosiin	H	OGlu	H	OH	H	H	H	OH	H	ditto
15	7 Apiin	H	OApq	H	OH	H	H	H	OH	H	ditto
	8 Luteolin	H	OH	H	OH	H	H	OH	OH	H	ditto
	9 Galuteolin	H	OH	H	OGlu	H	H	OH	OH	H	ditto
20	10 Gluco -luteolin	H	OGlu	H	OH	H	H	OH	OH	H	ditto
	11 Acacetin	H	OH	H	OH	H	H	H	OCH ₃	H	ditto
	12 Linarin	H	ORut	H	OH	H	H	H	OCH ₃	H	ditto
25	13 Diosmetin	H	OH	H	OH	H	H	OH	OCH ₃	H	ditto
	14 Diosmin	H	ORut	H	OH	H	H	OH	OCH ₃	H	ditto
	15 Baicalein	H	OH	OH	OH	H	H	H	H	H	ditto
30	16 Fisetin	H	OH	H	H	OH	H	H	OH	H	ditto
	17 Kaempferol	H	OH	H	OH	OH	H	H	OH	H	ditto
	18 Trifolin	H	OH	H	OH	OGal	H	H	OH	H	ditto
35	19 Astragalin	H	OH	H	OH	OGlu	H	H	OH	H	ditto

Table 1 (continued)

	No.	Name	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹	X
5	20	Robinin	H	ORha	H	OH	ORob	H	H	OH	H	Double
	21	Kaem -pferitrin	H	ORha	H	OH	ORha	H	H	OH	H	ditto
10	22	Quercetin	H	OH	H	OH	OH	H	OH	OH	H	ditto
	23	Quercitrin	H	OH	H	OH	ORha	H	OH	OH	H	ditto
	24	Iso -quercitrin	H	OH	H	OH	OGLu	H	OH	OH	H	ditto
15	25	Rutin	H	OH	H	OH	ORut	H	OH	OH	H	ditto
	26	Morin	H	OH	H	OH	OH	OH	H	OH	H	ditto
	27	Myricetin	H	OH	H	OH	OH	H	OH	OH	OH	ditto
	28	Myricitrin	H	OH	H	OH	ORha	H	OH	OH	OH	ditto
20	29	Datiscetin	H	OH	H	OH	OH	OH	H	H	H	ditto
	30	Quer -cetagetin	H	OH	OH	OH	OH	H	OH	OH	H	ditto
25	31	Quer -cetagitrin	H	OGLu	OH	OH	OH	H	OH	OH	H	ditto
	32	Rhamnetin	H	OCH ₃	H	OH	OH	H	OH	OH	H	ditto
	33	Iso -rhamnetin	H	OH	H	OH	OH	H	OCH ₃	OH	H	ditto
30	34	Pinocembrin	H	OH	H	OH	H	H	H	H	H	Single
	35	Naringenin	H	OH	H	OH	H	H	H	OH	H	ditto
	36	Salipurpin	H	OH	H	OGLu	H	H	H	OH	H	ditto
	37	Prunin	H	OGLu	H	OH	H	H	H	OH	H	ditto
35	38	Naringin	H	ORha	H	OH	H	H	H	OH	H	ditto
	39	Sakuranetin	H	OCH ₃	H	OH	H	H	H	OH	H	ditto
	40	Sakuranin	H	OCH ₃	H	OGLu	H	H	H	OH	H	ditto
40	41	Hesperetin	H	OH	H	OH	H	H	OH	OCH ₃	H	ditto
	42	Hesperidin	H	ORut	H	OH	H	H	OH	OCH ₃	H	ditto
	43	Eriodictyol	H	OH	H	OH	H	H	OH	OH	H	ditto
	44	Eriodictin	H	ORha	H	OH	H	H	OH	OH	H	ditto
45	45	Pinobanksin	H	OH	H	OH	OH	H	H	H	H	ditto
	46	Arom -adendrin	H	OH	H	OH	OH	H	H	OH	H	ditto
	47	Engelitin	H	OH	H	OH	ORha	H	H	OH	H	ditto
50	48	Fustin	H	OH	H	H	OH	H	OH	OH	H	ditto
	49	Taxifolin	H	OH	H	OH	OH	H	OH	OH	H	ditto

Table 1 (continued)

No.	Name	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸	R ⁹	X
50	Astilbin	H	OH	H	OH	ORha	H	OH	OH	H	Single
51	Ampelopsin	H	OH	H	OH	OH	H	OH	OH	OH	ditto

OGlu: Glucoside, OApG: Apioglucoside, ORut: Rutinoside,
OGal: Galactoside, ORha: Rhamnoside, ORob: Robinobioside

The compounds Nos. 34 to 51 in Table 1 include a single bond as X. Thus, the carbon atom in the 2-position or the carbon atoms in the 2- and 3-positions are asymmetrical, and there exist stereoisomers. It is known that pinocembrin includes (±) and (S) isomers; naringenin, sakuranetin, hesperetin and eriodictyol include (±), (R), and (S) isomers; pinobanksin includes (2R-trans) and (2S-trans) isomers; aromadendrin and fustin include trans-(±), (2R-trans), and (2S-trans) isomers; taxifolin and ampelopsin include trans-(±), (2R-trans), (2S-trans), and (2R-cis) isomers.

It is possible to use, as the flavonoids, compounds isolated and purified from naturally occurring plants or chemically synthesized. Many compounds described in Table 1 are commercially available. For example, it is possible to obtain flavone, apigenin, luteolin, acacetin, linarin, diosmetin, baicalein, fisetin, kaempferol, quercetin, hesperetin, and hesperidin from Funakoshi Co., Ltd., Tokyo.

Examples of the acute toxicity of the present substance are as follows: Mouse LD₅₀ of quercetin (oral administration): 160 mg/kg and mouse LD₅₀ of fisetin (intravenous injection): 180 mg/kg.

Further, no abnormalities were observed for a week after hesperetin was administered orally to BALB/c mice (female, seven weeks old) at the dose of 100 mg/kg. The same results were obtained where hesperidin, acacetin, diosmetin, apigenin, luteolin, or kaempferol was administered.

As a pharmacological effect, the present substance exhibits the function to inhibit destruction of chondrocyte matrix in chondrocyte culture (derived from cartilage of rabbit shoulder and knee joints) (see Example 1 as below).

Accordingly, the present substance is useful as a chondroprotective agent for treating various types of arthropathy accompanying the cartilage destruction of joints. Examples of such arthropathy are rheumatoid arthritis, osteoarthritis, periarthritis humeroscapularis, shoulder-arm-neck syndrome, lumbago, etc.

The chondroprotective agent containing the present substance as an active ingredient may be in the form of any conventional formulation. The chondroprotective agent may contain the present substance alone, or a mixture of the present substance with any pharmaceutically acceptable carrier or diluent. The chondroprotective agent may contain the active ingredient in an amount of 0.01 to 100 percent by weight, preferably 0.1 to 70 percent by weight.

The chondroprotective agent of the present invention may be administered orally or by some other routes.

The dose of the chondroprotective agent according to the present invention varies with the patient (animal or human), age, individual differences, state of illness, and the like. Generally speaking, however, when a human is treated, the dose of oral administration of the present substance is in the range of 0.1 to 500 mg/kg (body weight) per day, preferably 0.5 to 200 mg/kg (body weight), which is usually divided into 1 to 4 dosages in a day, although the dose outside the above range may sometimes be administered.

EXAMPLE

The present invention now will be further illustrated by, but is by no means limited to, the following Examples.

Example 1: Effect of Test Compounds on Proteoglycan Depletion in Chondrocyte Culture

(a) Preparation of Cultured Chondrocytes

5 The cartilages were sterilely extracted from the shoulder and knee joints of rabbits (New Zealand White Rabbits) (body weight of 1 to 1.5 kg). The cartilages were thoroughly washed with PBS (-) (free of Ca^{2+} , Mg^{2+}), Hanks' solution and 0.1% EDTA-PBS (-), and then cut into small segments (1 mm x 1 mm x 1 mm). After PBS (-) containing 0.1% EDTA was added, the segments were allowed to stand in an incubator of 37 °C for 30 minutes. Then, the segments were treated with a trypsin solution (0.25%) at 37 °C for one hour
 10 to remove the connective tissue adhered to the cartilage. After the supernatant had been removed, the cartilages were treated for about 2 to 2.5 hours in a Ham F-12 medium containing 10 % fetal bovine serum (FBS) and 0.2 % collagenase. Then, the collagenase solution was centrifuged (1500 r.p.m.), and the residual chondrocytes were washed twice with a Ham F-12 medium (chondrocyte culture medium) containing 10 % FBS. Finally, the resulting suspension was adjusted so that the chondrocytes were suspended in the
 15 concentration of 3×10^5 cells/ml in the chondrocyte culture medium. The chondrocytes were seeded in an amount of 1 ml/well on 24-well plates. The chondrocytes became confluent after 4 days. The experiment were performed within two weeks after reaching the confluent stage.

(b) Addition of Compounds to be tested and Proteoglycan Depleting Agents

20 The chondrocyte culture medium which had been used for cultivating the chondrocytes was removed from each well and 800 μl of fresh serum-free S-Clone medium containing 0.1% human serum albumin was added. Further, 100 μl of S-Clone medium containing the compounds to be tested (containing the compound in the concentration of 10 fold the final concentration; DMSO concentration = 2.5%) was added.
 25 The chondrocytes were cultured in the presence of carbon dioxide (5%) and air (95%) for 2 hours. Then, the proteoglycan depleting agent, PMA (phorbol myristate acetate) (final concentration = 0.1 $\mu\text{g/ml}$) was added into the culture medium of the chondrocytes.

The compounds to be tested were as follows:

Compounds of present invention: apigenin (present substance No. 5), luteolin (present substance No. 8), acacetin (present substance No. 11), linarin (present substance No. 12), diosmetin (present substance No. 13), baicalein (present substance No. 15), fisetin (present substance No. 16), kaempferol (present substance No. 17), quercetin (present substance No. 22), hesperetin (present substance No. 41, (S) isomer), and hesperidin (present substance No. 42, (S) isomer) (all from Funakoshi Co.)

Comparative substance: Indometacin (Sigma Chemical Co.)

(c) Determination of proteoglycan

Proteoglycan depletion was determined by the measurement of the glycosaminoglycan (major constituent of proteoglycan, hereinafter referred to as GAG) content following digestion of the chondrocyte matrix
 40 with papain.

After 2 days, the supernatant of the chondrocyte culture was removed. Then, 1 ml of 0.03% papaine solution was added to the remaining chondrocyte matrix layer and a reaction was performed at 65 °C for 1 hour to liberate the GAG from the matrix layer. The content of the GAG in the treated papaine solution was determined by the 1,9-dimethylmethylene blue method (refer to R.W. Farndale, Biochim. Biophys. Acta.,
 45 Vol. 883, pp. 173 to 177, 1986). The GAG content in the chondrocyte matrix of the control test wherein the proteoglycan depleting agent was not added was shown as "100", and the relative amount of the GAG of each experiment except the control test was calculated by the following formula:

$$\text{GAG relative amount (\%)} = (\text{B/A}) \times 100$$

50 wherein A represents the GAG content of the control tests wherein the proteoglycan depleting agent was not added, and B represents the GAG content wherein the proteoglycan depleting agents were added alone or the GAG content wherein the proteoglycan depleting agents and the compounds to be tested were added.

55 The GAG contents of the control tests varied in a range of 11.23 to 59.0 $\mu\text{g/ml}$, depending on the period from the time when the chondrocytes became confluent until the time when the chondrocytes were used in the above experiment.

The results are shown in Table 2. The GAG content is the value of the mean value \pm standard error ($n = 3$ to 6). For each of the compounds to be tested, the control test and the proteoglycan depleting test wherein the proteoglycan depleting agent was added were carried out and the results thereof are also shown. The significance was determined by Student's t-test with respect to the proteoglycan depleting test wherein the proteoglycan depleting agent was added. The results of the determination are shown as follows:

*: $P < 0.05$;

**: $P < 0.01$;

***: $P < 0.001$.

In comparison with the GAG content in the control tests wherein the proteoglycan depleting agent was not added, the addition of the proteoglycan depleting agents, PMA, induced a loss of GAG content. Under these conditions, the present compound significantly inhibited or reduced the loss of GAG content, and showed a function to inhibit or suppress the proteoglycan depletion. On the other hand, indomethacin, a conventional analgesic and anti-inflammatory agent, did not show the function to inhibit or suppress the proteoglycan depletion, but caused a significant exacerbation on the proteoglycan depletion.

Table 2

Samples	GAG content ($\mu\text{g/ml}$)	(Relative amount of GAG) (%)
Control	54.7 \pm 0.8***	(100)
PMA	16.5 \pm 0.7	(30.2)
PMA + No. 5 (100 μM)	33.3 \pm 0.7***	(60.9)
Control	54.8 \pm 0.5***	(100)
PMA	15.2 \pm 0.6	(27.7)
PMA + No. 8 (100 μM)	28.6 \pm 0.5***	(52.2)
PMA + No. 17 (100 μM)	30.5 \pm 0.3***	(55.7)
Control	54.0 \pm 1.2***	(100)
PMA	20.0 \pm 0.4	(37.0)
PMA + No. 11 (100 μM)	24.2 \pm 0.3	(44.8)
PMA + No. 16 (100 μM)	30.1 \pm 0.9***	(55.7)
Control	55.3 \pm 0.6***	(100)
PMA	17.5 \pm 0.7	(31.6)
PMA + No. 12 (100 μM)	19.8 \pm 0.7*	(35.8)
PMA + No. 13 (100 μM)	27.1 \pm 0.7***	(49.0)
Control	11.23 \pm 0.2***	(100)
PMA	4.94 \pm 0.1	(44.0)
PMA + No. 15 (100 μM)	7.67 \pm 0.5*	(68.3)
Control	56.1 \pm 0.8***	(100)
PMA	20.4 \pm 0.7	(36.4)
PMA + No. 22 (100 μM)	31.0 \pm 0.6***	(55.3)
PMA + No. 42 (100 μM)	24.0 \pm 0.6**	(42.8)
Control	59.0 \pm 0.9***	(100)
PMA	21.1 \pm 0.6	(35.8)
PMA + No. 41 (100 μM)	28.7 \pm 0.4***	(48.6)
Control	28.0 \pm 0.7***	(100)
PMA	15.4 \pm 0.5	(55.0)
PMA + indometacin (10 μM)	13.2 \pm 0.6*	(47.1)
(33 μM)	11.7 \pm 0.8**	(41.8)

Example 2: Formulation of Granule

The following ingredients were mixed homogeneously:

Apigenin	20 parts by weight
Lactose	68 parts by weight
Low-substituted hydroxypropylcellulose	10 parts by weight
Hydroxypropylcellulose	2 parts by weight

The mixture was kneaded using 32 parts by weight of a wetting agent, ethanol. Then, the kneaded mixture was granulated by wet granulation and dried to obtain the granule.


As explained above, the present substance strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage. Further, the present substance has low toxicity.

Accordingly, the present substance is very useful for the treatment of arthropathy, such as rheumatoid arthritis, osteoarthritis, periartthritis humeroscapularis, shoulder-arm-neck syndrome, lumbago, and so on.

Although the present invention has been described with reference to specific embodiments, various changes and modifications obvious to those skilled in the art are deemed to be within the spirit, scope, and concept of the invention.

Claims

1. A chondroprotective agent comprising a flavonoid compound of the general formula (I):



wherein R¹ to R⁹ are, independently, a hydrogen atom, hydroxyl group, or methoxyl group and X is a single bond or a double bond, or a stereoisomer thereof, or a naturally occurring glycoside thereof, and a pharmaceutically acceptable carrier.

2. A chondroprotective agent according to claim 1, wherein X is a double bond.
3. A chondroprotective agent according to claim 2, wherein said flavonoid compound is one or more compounds selected from the group consisting of flavone, chrysin, toringin, primetin, apigenin, cosmosiin, apiin, luteolin, galuteolin, glucoluteolin, acacetin, linarin, diosmetin, diosmin, baicalein, fisetin, kaempferol, trifolin, astragalin, robinin, kaempferitrin, quercetin, quercitrin, isoquercitrin, rutin, morin, myricetin, myricitrin, datiscetin, quercetagetin, quercetagitrin, rhamnetin, and isorhamnetin.
4. A chondroprotective agent according to claim 1, wherein X is a single bond.
5. A chondroprotective agent according to claim 4, wherein said flavonoid compound is one or more compounds selected from the group consisting of pinocembrin, naringenin, salipurpin, prunin, naringin, sakuranetin, sakuranin, hesperetin, hesperidin, eriodictyol, eriodictin, pinobanksin, aromadendrin, engelitin, fustin, taxifolin, astilbin, and ampelopsin.
6. A chondroprotective agent according to claim 1, wherein said naturally occurring glycoside is one or more compounds selected from the group consisting of glucoside, galactoside, fructoside, rhamnoside, rutinoside, arabinoside, xyloside, apioglucoside, and robinobioside.

7. A chondroprotective agent according to claim 1, wherein said flavonoid compound is one or more compounds selected from the group consisting of apigenin, luteolin, acacetin, linarin, diosmetin, baicalein, fisetin, keampferol, quercetin, hesperetin (S-form), and hesperidin (S-form).
- 5 8. A chondroprotective agent according to claim 1, wherein said flavonoid compound, stereoisomer thereof, or naturally occurring glycoside thereof is an extract from a naturally occurring material.
9. Use of a compound of the general formula (I), or a stereoisomer thereof, or a naturally occurring glycoside thereof according to claim 1, for preparing a chondroprotective agent.

10

15

20

25

30

35

40

45

50

55

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 633 022 A3

(12)

EUROPEAN PATENT APPLICATION(21) Application number: **94109872.5**(51) Int. Cl.⁶: **A61K 31/365, A61K 31/70**(22) Date of filing: **27.06.94**(30) Priority: **09.07.93 JP 194182/93**(43) Date of publication of application:
11.01.95 Bulletin 95/02(64) Designated Contracting States:
CH DE FR GB IT LI SE(68) Date of deferred publication of the search report:
02.08.95 Bulletin 95/31

(71) Applicant: **KUREHA CHEMICAL INDUSTRY CO., LTD.**
9-11, Horidome-cho, 1-chome
Nihonbashi
Chuo-ku
Tokyo 103 (JP)


(72) Inventor: **Watanabe, Koju**

2-5-7, Tsurumai
Sakado-shi,
Saitama 350-02 (JP)
Inventor: **Nilimura, Koichi**
Rune Warabi 1-718,
1-17-30, Chuo
Warabi-shi,
Saitama 335 (JP)
Inventor: **Umekawa, Kiyonori**
5-4-309, Mihama
Urayasu-shi,
Chiba 279 (JP)

(74) Representative: **Minderop, Ralph H., Dr. rer.**
nat. et al
Cohausz & Florack
Patentanwälte
Postfach 33 02 29
D-40435 Düsseldorf (DE)

(54) **Chondroprotective agents.**

(57) A chondroprotective agent comprising a flavonoid compound of the general formula (I):



wherein R¹ to R⁹ are, independently, a hydrogen atom, hydroxyl group, or methoxyl group and X is a single bond or a double bond, or a stereoisomer thereof, or a naturally occurring glycoside thereof is disclosed. The above compound strongly inhibits proteoglycan depletion from the chondrocyte matrix and exhibits a function to protect cartilage, and thus, is extremely effective for the treatment of arthropathy.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 10 9872

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	AGENTS AND ACTIONS, vol. 12, page 298 O.P. AGARWAL 'The anti-inflammatory action of nepitrin, a flavonoid' * See the whole document, especially Table 2 and penultimate paragraph of the discussion *	1,2,9	A61K31/365 A61K31/70
Y	* " *	1-3,6-9	
Y	--- Z. RHEUMATOL., vol. 42, page 203 G. WILHELMI ET AL. 'zur Prüfung potentieller Antiarthrotika an der spontanen Arthrose der Maus' *see the whole document *	1-3,6-9	
X	--- CHEM. PHARM. BULL., vol. 32, page 2724 M. KUBO ET AL. 'Studies on scutellariae radix. VII Anti-arthritic and anti-inflammatory actions of methanolic extract and flavonoid components from scutellariae radix' * see the whole document, especially Table 4 *	1-3,6-8	TECHNICAL FIELDS SEARCHED (Int.Cl.6) A61K
Y	* " *	9	
Y	--- US-A-4 268 517 (NIEBES ET AL.) 19 May 1981 * see the whole document* --- -/--	1-3,6-9	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 15 May 1995	Examiner Isert, B
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background U : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 10 9872

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	ARCH. PHARM. RES., vol. 16, no. 1, 1993 pages 25-28, S.J. LEE ET AL. 'antiinflammatory activity of naturally occurring flavone and flavonol glycosides' * see the whole document, especially Figure 1 and Table 1 *	1-4, 6-8	
Y	* " "	9	
X	--- FITOTERAPIA , vol. 90, no. 5, 1990 pages 460-461, G.B.S. REDDY ET AL. 'comparison of the anti-inflammatory activity of certain methoxyflavonoids' * see the summary *	1-3, 6-8	
Y	* " *	9	
X, P	--- INDIAN J. PHARMACOL. SCI., vol. 55, no. 5, September 1993 pages 180-183, N.S. MUTHIAH ET AL. 'Anti-inflammatory activity of flavone and its methoxy derivatives- a structure activity study-' * see the summary and Table 2 *	1-4, 6-8	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X, P	* " *	9	
A	--- EP-A-0 395 441 (KUREHA KAGAKU KOGYO) 31 October 1990 * see page 8, line 36 - page 9, line 30; claims 1-8, 10 *	1-9	

The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 15 May 1995	Examiner Isert, B
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			



European Patent
Office

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions.

namely:

See sheet -B-

- ☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent
Office

EP 94 10 9872 -B-

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims 1 (partially), 2,3,6-9 (partially):
Chondroprotective flavones
2. Claims 1 (partially), 4,5,6-9 (partially):
Chondroprotective flavanones

The first part of the document discusses the importance of maintaining accurate records of all transactions. It emphasizes that every entry, no matter how small, should be carefully documented to ensure the integrity of the financial data. This includes recording dates, amounts, and the nature of the transactions. The second part of the document provides a detailed overview of the accounting process, from initial data collection to the final reporting stage. It outlines the various steps involved in reconciling accounts, identifying discrepancies, and preparing financial statements. The third part of the document focuses on the analysis of the financial data, highlighting key trends and potential areas of concern. It discusses the importance of comparing current performance with historical data and industry benchmarks. The final part of the document offers recommendations for improving financial management practices, such as implementing stricter controls and enhancing communication between departments. Throughout the document, the emphasis is on transparency, accuracy, and proactive management of financial resources.